Catalogue of American Amphibians and Reptiles.

Lovich, J.E. and C.J. McCoy. 1994. *Graptemys gibbonsi.*

Graptemys gibbonsi Lovich and McCoy Pascagoula Map Turtle

Graptemys pulchra: Cagle, 1952:223 (part).
Graptemys gibbonsi Lovich and McCoy, 1992:293. Type-locality, "Chickasawhay River, Leakesville, Greene County, Mississippi, USA." Holotype, Carnegie Museum 94979, an adult male collected by R.C. Vogt, M. Pappas, and P.S. Freed on 21 July 1978.

• **Content.** No subspecies are recognized.

• **Definition.** The Pascagoula Map Turtle is a large (to 295 mm carapace length) species. The shell is high-domed, possesses a median keel, and has a single vertical yellow bar on the dorsal surface of each marginal scute. The median carapace keel is composed of prominent spines on the posterior portions of the second and third vertebrals. A complete or broken black stripe, most pronounced anteriorly, marks the median keel of the vertebrals. Pleural scutes 1-3 have circular yellow markings on the distal parts. The plastron is pale yellow with dark pigment on some seams. Ground color of the head and limbs is brown to olive with light yellow or yellowish-green stripes and blotches. The head pattern consists of a large interorbital blotch that is connected to large postorbital blotches on either side of the head. Supraoccipital spots are absent. The dorsal paramedian neck stripes are not expanded anteriorly, but may contact the postorbital blotches. A three-pronged yellow blotch (nasal trident) is usually present on the dorsal head surface behind the nares.

Adult females can be well over twice the length and weight of adult males (Gibbons and Lovich, 1990), and have conspicuously enlarged heads with broad jaw surfaces. Males have longer tails with the vent posterior to the rim of the carapace. Both sexes have relatively flat plastra.

• **Descriptions.** Complete descriptions are in Lovich and McCoy (1992), McCoy and Lovich (in press), and Ernst et al. (1994). Prior descriptions are clouded by the fact that the Pascagoula Map Turtle was formerly considered to be *G. pulchra* (in part). The reader is referred to Lovich (1985) for these composite descriptions.

• **Illustrations.** Color photographs of *Graptemys gibbonsi* are found in Ashton and Ashton (1985), Ernst and Barbour (1972), and Ernst et al. (1994). Black and white photographs are in Ernst and Barbour (1972, 1989), Mount (1975), and Wahlquist (1970). Composite illustrations of *G. gibbonsi* (as *G. pulchra*, in part) are listed in Lovich and McCoy (1992).



Map. The range of *Graptemys gibbonsi*. The large circle marks the type-locality, dots indicate other selected localities (from Iverson, 1992).

• **Distribution.** *Graptemys gibbonsi* is found only in large to medium-sized rivers in the Pascagoula River and Pearl River systems, Mississippi and eastern Louisiana. Pascagoula River system localities are in the Pascagoula, Leaf, and Chickasawhay rivers; Red, Bowie, and Okatoma creeks (Cliburn, 1971); and Tallahala Creek (J.L. Dobie, pers. comm.). *Graptemys gibbonsi* does not occur in either Big Creek or the Escatawpa River, which are Alabama tributaries of the Pascagoula River (Mount, 1975), or in Black Creek, a major tributary of the Pascagoula in Mississippi (Cliburn, 1971). Pearl River drainage populations occur in Ross Barnett Reservoir (Boyd and Vickers, 1963), and in the Pearl and Bogue Chitto rivers (Lovich and McCoy, 1992). Dundee and Rossman (1989) published a record of this species (as *G. pulchra*) from the Tickfaw River at U.S. Highway 190,



Figure. A male Graptemys gibbonsi from the Chickasawhay River in Greene County, Mississippi. Photograph by J.E. Lovich.

Livingston Parish, Louisiana. Lovich and McCoy (1992) questioned that record, as the species is unknown in the Tangipahoa, Tchefuncte, or Amite rivers, which lie between the Tickfaw River and Bogue Chitto River (Cagle, 1952; Cliburn, 1971).

• **Fossil Record.** None.

• **Pertinent Literature.** Literature prior to the recognition of *Graptemys gibbonsi* as a separate species in the *Graptemys pulchra* complex is summarized in Lovich (1985). Tinkle (1962), Little (1973), and Lovich and Ernst (1989) examined scute morphology in *G. pulchra* (*sensu lato*), including specimens of *G. gibbonsi*. Literature dealing specifically with turtle populations now recognized as *G. gibbonsi* is listed by topic: collecting techniques (Chaney and Smith, 1950), hatchling behavior (Anderson, 1958), natural history data on reproduction, diet, habitat, and conservation (Cagle, 1952; Dundee and Rossman, 1989; McCoy and Lovich, in press; Ernst et al., 1994), geographic distribution (Boyd and Vickers, 1963; Cliburn, 1971), sexual size dimorphism (Gibbons and Lovich, 1990), blood proteins (McKown, 1972), and systematics and taxonomy (Lovich and McCoy, 1992; Lamb et al., 1994).

• **Etymology.** The specific epithet *gibbonsi* is a patronym honoring Dr. J. Whitfield Gibbons of the University of Georgia, Savannah River Ecology Laboratory.

• **Comment.** The Pascagoula Map Turtle was described by Lovich and McCoy (1992), who demonstrated that the species *Graptemys pulchra* Baur (1893:675) is a composite consisting of three distinct, allopatric species. The other species in the complex include *G. pulchra* (*sensu stricto*) of rivers tributary to the Mobile Bay Drainage in Alabama, Mississippi, and northwestern Georgia; and *G. ernsti* in the Escambia Bay drainage of southeastern Alabama and western Florida. The three species are well-differentiated on the basis of morphology and color pattern, and exhibit divergent mtDNA genotypes. These three species, together with *G. barbouri*, form the *Graptemys pulchra* species group (Lovich and McCoy, 1992; Lamb et al., 1994). Additional information on *G. gibbonsi* (as *G. pulchra*, in part) was summarized by Lovich (1985).

Literature Cited

Anderson, P.K. 1958. The photic response and water-approach behavior of hatchling turtles. Copeia 1958:211-215.

Ashton, R.E., Jr. and P.S. Ashton. 1985. Handbook of the reptiles and amphibians of Florida. Part II. Lizards, turtles and crocodilians. Windward Publishing Co., Miami, Florida.

Baur, G. 1893. Two new species of North American Testudinata. Amer. Nat. 27:675-677.

Boyd, C.E., and D.H. Vickers. 1963. Distribution of some Mississippi amphibians and reptiles. Herpetologica 19:202-205.

Cagle, F.R. 1952. The status of the turtles *Graptemys pulchra* Baur and *Graptemys barbouri* Carr and Marchand, with notes on their natural history. Copeia 1952:223-234.

Chaney, A. and C.L. Smith. 1950. Methods for collecting map turtles. Copeia 1950:323-324.

Cliburn, J.W. 1971. The ranges of four species of *Graptemys* in Mississippi. J. Mississippi Acad. Sci. 16:16-19.

Dundee, H.A. and D.A. Rossman. 1989. The amphibians and reptiles of Louisiana. Louisiana State Univ. Press, Baton Rouge.

Ernst, C.H. and R.W. Barbour. 1972. Turtles of the United States. Univ. Press Kentucky, Lexington.

—— and ——. 1989. Turtles of the world. Smithsonian Institution Press, Washington, D.C.

——, J.E. Lovich, and R.W. Barbour. 1994. Turtles of the United States and Canada. Smithsonian Institution Press, Washington, D.C.

Gibbons, J.W. and J.E. Lovich. 1990. Sexual dimorphism in turtles with emphasis on the slider turtle (*Trachemys scripta*). Herpetol. Monogr. 4:1-29.

Iverson, J.B. 1992. A revised checklist with distribution maps of the turtles of the world. Privately Printed, Richmond, Indiana.

Lamb, T., C. Lydeard, R.B. Walker, and J.W. Gibbons. 1994. Molecular systematics of map turtles (*Graptemys*): a comparison of mitochondrial restriction site versus sequence data. Syst. Biol. 43:543-559.

Little, R.B. 1973. Variation in the plastral scutellation of *Graptemys pulchra* (Reptilia, Chelonia, Emydidae). A.S.B. Bull. 20:65-66.

Lovich, J.E. 1985. *Graptemys pulchra* Baur. Alabama map turtle. Cat. Amer. Amphib. Rept.:360.1-360.2.

—— and C.H. Ernst. 1989. Variation in the plastral formulae of selected turtles with comments on taxonomic utility. Copeia 1989:304-318.

—— and C.J. McCoy. 1992. Review of the *Graptemys pulchra* group (Reptilia, Testudines, Emydidae), with descriptions of two new species. Ann. Carnegie Mus. 61:293-315.

McCoy, C.J. and J.E. Lovich. In press. *Graptemys gibbonsi*, Pascagoula Map Turtle. *In* P.C.H. Pritchard and A. Rhodin (eds.), Conservation of freshwater turtles. IUCN, Species Survival Commission.

McKown, R.R. 1972. Phylogenetic relationships within the turtle genera *Graptemys* and *Malaclemys*. Ph.D. diss., Univ. Texas, Austin.

Mount, R.H. 1975. The reptiles and amphibians of Alabama. Auburn Univ. Agr. Exp. Sta., Auburn, Alabama.

Tinkle, D.W. 1962. Variation in shell morphology of North American turtles. I. The carapacial seam arrangements. Tulane Stud. Zool. 9:331-349.

Wahlquist, H. 1970. Sawbacks of the Gulf Coast. Internatl. Turtle Tortoise Soc. J. 4:10-13, 28.

Jeffrey E. Lovich, United States Department of the Interior, National Biological Survey, Palm Springs Field Station, Midcontinent Ecological Science Center, 63-500 Garnet Avenue, P.O. Box 2000, North Palm Springs, CA 92258-2000, and **Clarence J. McCoy**, deceased.

Primary editor for this account, Michael E. Seidel.

Published 30 December 1994 and Copyright © 1994 by the Society for the Study of Amphibians and Reptiles.