

## Geology

### Mantle plume influence on the Neogene uplift and extension of the U.S. western Cordillera?

Tom Parsons, George A. Thompson and Norman H. Sleep

*Geology* 1994;22:83-86

doi: 10.1130/0091-7613(1994)022<0083:MPIOTN>2.3.CO;2

---

#### Email alerting services

click [www.gsapubs.org/cgi/alerts](http://www.gsapubs.org/cgi/alerts) to receive free e-mail alerts when new articles cite this article

#### Subscribe

click [www.gsapubs.org/subscriptions/](http://www.gsapubs.org/subscriptions/) to subscribe to *Geology*

#### Permission request

click <http://www.geosociety.org/pubs/copyrt.htm#gsa> to contact GSA

Copyright not claimed on content prepared wholly by U.S. government employees within scope of their employment. Individual scientists are hereby granted permission, without fees or further requests to GSA, to use a single figure, a single table, and/or a brief paragraph of text in subsequent works and to make unlimited copies of items in GSA's journals for noncommercial use in classrooms to further education and science. This file may not be posted to any Web site, but authors may post the abstracts only of their articles on their own or their organization's Web site providing the posting includes a reference to the article's full citation. GSA provides this and other forums for the presentation of diverse opinions and positions by scientists worldwide, regardless of their race, citizenship, gender, religion, or political viewpoint. Opinions presented in this publication do not reflect official positions of the Society.

---

#### Notes



# Mantle plume influence on the Neogene uplift and extension of the U.S. western Cordillera?

Tom Parsons

U.S. Geological Survey, MS 999, 345 Middlefield Road, Menlo Park, California 94025

George A. Thompson

Norman H. Sleep

Department of Geophysics, Stanford University, Stanford, California 94305

## ABSTRACT

Despite its highly extended and thinned crust, much of the western Cordillera in the United States is elevated more than 1 km above sea level. Therefore, this region cannot be thought of as thick crust floating isostatically in a uniform mantle; rather, the lithospheric mantle and/or the upper asthenosphere must vary in thickness or density across the region. Utilizing crustal thickness and density constraints, we modeled the residual mass deficit that must occur in the mantle lithosphere and asthenosphere beneath the western Cordillera. A major hot spot broke out during a complex series of Cenozoic tectonic events that included lithospheric thickening, back-arc extension, and transition from a subduction to a transform plate boundary. We suggest that many of the characteristics that make the western Cordillera unique among extensional provinces can be attributed to the mantle plume that created the Yellowstone hot spot.

## INTRODUCTION

Drawing on the growing understanding of mantle plumes and associated flood basalts (e.g., Richards et al., 1989; Sleep, 1990; Griffiths and Campbell, 1991; Hill et al., 1992) leads to the conclusion that the Yellowstone plume strongly influenced late Cenozoic tectonic events in the United States western Cordillera. Here we explore the probable consequences of the emergence of the hot spot about 17 Ma in northwestern Nevada, which was accompanied by voluminous basaltic volcanism on the Columbia Plateau and propagation of the northern Nevada rift hundreds of kilometres southward (e.g., Zoback and Thompson, 1978; Pierce and Morgan, 1992). Unlike many other highly extended terranes, the northern Basin and Range province is elevated to an average 1.5 km above sea level. The western Cordillera crust is generally not thick enough to support its elevation isostatically, which implies that a deficit in mantle density has caused the uplift (e.g., Eaton, 1982; Morgan and Swanberg, 1985). The northern Basin and Range is the broadest part of the province, which began in an extensional back-arc setting during Tertiary subduction off the west coast of North America (e.g., Eaton, 1979; Zoback et al., 1981). The total amount of crustal extension across the Basin and Range province estimated from surface geology is ~50%–300% (e.g., Hamilton and Myers, 1966; Zoback et al., 1981; Wernicke, 1992).

## MANTLE PLUMES

Whereas the deep origin of mantle plumes is controversial, their surface expression is dramatic and widely observed. Perhaps the two most ubiquitous manifestations of mantle plumes are voluminous volcanism and a broad regional topographic swell. The typical swell associated with plumes is ~1–2 km of uplift centered across a region ~1000–2000 km in diameter (e.g., Crough, 1983; Sleep, 1990), which is the result of isostatic compensation in the asthenosphere and lithosphere and, to a lesser extent, a dynamic pressure gradient of flow in the asthenosphere (below the limit of detection at Hawaii) (Sleep, 1990).


The interpreted anatomy of a mantle plume consists of a starting plume head generated during its initial ascent through the mantle, an active plume tail that continues to flow after the starting plume

head contacts the lithosphere, and ponded plume-tail material that collects as the active plume tail continues to flow (e.g., Griffiths and Campbell, 1991). The initial contact of the starting plume head with the lithosphere heats a broad region because the plume head is much wider than its tail (e.g., Campbell and Griffiths, 1990; Duncan and Richards, 1991). When a hot-spot swell forms in continental lithosphere the uplifted crust may be raised far enough above the level of mid-plate compression to be in a state of extension, which results in rift formation centered in the swell (e.g., Crough, 1983). The magnitudes of the generated deviatoric stresses fall short of those necessary to cause complete continental breakup (Hill, 1991).

The buoyancy and swell associated with the starting plume head tend to move with the lithospheric plate (Sleep, 1990; Griffiths and Campbell, 1991). The heat associated with mantle plumes tends to be contained in the mantle near the plume for a considerable time. For example, Davies (1992) concluded that the thermal anomaly associated with the Hawaiian plume has not waned during its 43 m.y. existence because the swell does not decay monotonically with distance along the volcanic chain, as would be the case in a thermal decline with age. That is, the material hotter than the normal mantle adiabatic level remains ponded below the lithosphere and above normal asthenosphere because of the long time required for heat applied at the base of the lithosphere to conduct to the surface and because secondary convection within the thermal boundary created by plume material is inefficient.

## Yellowstone Plume

During the past 16–17 m.y. the North American plate has moved southwest over the Yellowstone plume, leaving the Snake River Plain behind as its track (Morgan, 1972; Armstrong et al., 1975; Pierce and Morgan, 1992). The Yellowstone hot spot emerged with a burst of basaltic volcanism that formed the Columbia Plateau flood basalts to the northwest and the northern Nevada rift to the southeast (Zoback and Thompson, 1978). Magmatic activity on the eastern Snake River Plain was characterized initially by pulses of silicic volcanism (each of ~2–3 m.y. duration) progressing toward the present Yellowstone caldera. Most of the silicic volcanic rocks decrease in age to the northeast (35 to 40 mm/yr) as a result of



**Figure 1.** Color contour plot of relative variation in upper-mantle mass beneath investigated part of western Cordillera. Values were calculated using modified form of isostatic equation with crustal-depth information from seismic-refraction profiles (shown as gray lines). Free-air gravity anomaly across region is near zero, which implies that area is in isostatic equilibrium. Thin crust of northern Basin and Range province associated with its high elevation requires underlying low-density upper mantle if isostasy is to be satisfied. We suggest that Yellowstone plume has supplied hot, low-density material to base of lithosphere, causing broad mantle anomaly pictured.

southwestward plate migration over the plume (Christiansen and Lipman, 1972; Armstrong et al., 1975). Extensive basaltic volcanism followed, covering the plain and persisting through Holocene time (e.g., Luedke and Smith, 1983). Seismic-refraction data indicate that much of the mid-crust beneath the plain was replaced by intruded basalt (e.g., Sparlin et al., 1982). The location of the Yellowstone starting plume head and the start of the plume tail is subject to discussion because the plume may have encountered the subducting Juan de Fuca plate during its ascent, and part of the plume-head material could have been spread westward as far as the Juan de Fuca Ridge by the descending plate (Hill et al., 1992).

#### WESTERN CORDILLERA MANTLE

Because of its relatively thin crust and high elevation, the western Cordillera must depend for isostatic support on mantle material of lower relative density beneath it. We compiled crustal thickness information for the western United States from seismic refraction surveys (Warren, 1969; Prodehl, 1978; Catchings and Mooney, 1989; Kohler and McCarthy, 1990; McCarthy et al., 1991) (Fig. 1) and corrected for upper-crustal density variation by using a basin-stripped isostatic residual gravity map (Saltus, 1991). These gravity data give an indication of the relative density variation of the upper-crustal basement rocks because the effects of sedimentary basins and regional trends have been removed. Density perturbations of the upper-crustal rocks can be approximated with the relation  $\rho \approx (g/0.04L_{uc})$  (Simpson and Jachens, 1989), where  $g$  is gravity in mgal and  $L_{uc}$  is an assumed upper-crustal thickness (15 km). Lower-crustal

densities are not delimited by the data and were assigned a uniform value of  $2.9 \text{ g/cm}^3$ ; the lower-crustal layer is too thin for even strong density variations within it to cause the large observed mass deficit. Combining the crustal-thickness and upper-crustal density data results in the approximate crustal contribution to uplift across the western Cordillera; this, when compared with the topography, indicates the mantle contribution to uplift (Fig. 1). We assume regional isostasy because the free-air gravity anomaly averages zero across the region (e.g., Thompson and Zoback, 1979).

We find the upper-mantle mass per unit area by dividing the study area into a grid of isostatic columns. These columns are calculated using a two-layer isostatic expression with a known crustal layer, and a second mantle layer that extends down to an assumed local isodensity asthenosphere at a depth of  $\sim 200$  km. The problem can be viewed as a group of 200-km-deep columns each containing the unknown boundary between the lithosphere (of unknown density) and the fixed-density asthenosphere. Because we treat the crust as a known quantity and we know the topographic elevation, the mass contained within the lithospheric mantle part of the column required to float it to its individual height can be calculated. The mantle mass per unit area of a given column ( $\Delta M_n$ ) is found by comparing it to a reference column, which is the column in the study area that contains the greatest amount of mass (and hence has the lowest mantle buoyancy):


$$\Delta M_n = (I_n - I_{ref}),$$

where  $I_n = \rho_a E_n + (\rho_a - \rho_c) L_{cn}$ , is a modified form of the isostatic equation (e.g., Lachenbruch and Morgan, 1990). The variable  $L_{cn}$  is the thickness of the crust in column  $n$ ,  $\rho$  is density, the  $a$  and  $c$  subscripts denoting asthenosphere and crust, respectively ( $\rho_a = 3.2 \text{ g/cm}^3$ ; variation in the fixed asthenosphere density does not affect relative difference between columns), and  $E_n$  is topographic elevation.

On the basis of isostatic calculations, Figure 1 shows that a broad low-density anomaly underlies the northern Basin and Range province, extending from the Sierra Nevada in the west to the Colorado Plateau in the east. The peak of the anomaly corresponds approximately to the inferred location of the initial breakout of the Yellowstone plume at McDermitt caldera (Zoback and Thompson, 1978); the bulk of the anomaly is to the south, beneath the central Basin and Range province. The actual plume track (eastern Snake River Plain) does not appear as a low-density anomaly on this image because the crust there has been largely replaced by more dense basalt (e.g., Sparlin et al., 1982), which has caused subsidence.

We calculate that the sum of the mantle isostatic mass deficit is  $\sim 3.6 \times 10^{18} \text{ kg}$  beneath the region investigated. Extension and thinning of the Basin and Range lithosphere allowed the asthenosphere to rise and take its place, causing some of the mass deficit. Earthquake-source methods that measure the variation in lithospheric thickness between the Basin and Range province and the unextended Colorado Plateau converge on a difference in the mantle-lid thickness that ranges between 15 and 50 km across the two provinces (Iyer and Hitchcock, 1989; Beghoul et al., 1993). If a reasonable density contrast between asthenosphere and lithosphere of  $0.05 \text{ g/cm}^3$  is used (e.g., Thompson and Zoback, 1979; Lachenbruch and Morgan, 1990), then the mass deficit caused by lithospheric thinning amounts to  $\sim 4.8 \times 10^{17}$ – $1.6 \times 10^{18} \text{ kg}$ , or 13%–44% of the total anomaly. If the entire anomaly is attributed to lithospheric thinning over an adiabatic asthenosphere, then a mantle lithospheric layer  $\sim 112$  km thick would have to have been removed from the Basin and Range province across the study area to account for the entire observed mass deficit.

**Figure 2. Schematic Tertiary tectonic history of western Cordillera. A: Horizontal subduction of Farallon plate thickens Cordillera lithosphere. B: Transition to normal-angled subduction creates back-arc extension in Basin and Range and thins lithosphere there. C: Yellowstone plume contacts lithosphere, ponding beneath thinnest regions. Northern Basin and Range province is centered above plume and thus is peak of topographic swell. Excess elevation triggers new phase of large-magnitude extension in Basin and Range which thins lithosphere relative to surrounding provinces. D: Subduction terminates, and extension continues in Basin and Range province, causing subsidence relative to surrounding regions.**




The calculated buoyancy flux of the Yellowstone plume is  $\sim 1.5 \times 10^3 \text{ kg s}^{-1}$  (Sleep, 1990). Thus the active Yellowstone plume tail could have generated  $\sim 20\%$  ( $\sim 7.6 \times 10^{17} \text{ kg}$ ) of the total mass deficit as ponded plume-tail material during the time (16–17 m.y.) it took to track from northern Nevada to its current position at Yellowstone. The time required for the Yellowstone plume tail to supply the mantle mass deficit not caused by lithospheric thinning ( $\sim 2.0 \times 10^{18}$ – $3.1 \times 10^{18} \text{ kg}$ ) ranges from 42 to 89 m.y., which is much longer than the age of the Yellowstone plume. Because the Yellowstone plume is probably about 16–17 m.y. old, we suggest that the plume had a starting head, the remnant of which causes an  $\sim 1.2 \times 10^{18}$ – $2.4 \times 10^{18} \text{ kg}$  mass deficit beneath the investigated part of the western Cordillera. If the plume head was roughly symmetric, then these numbers extrapolate to a total mass deficit of about  $1.7 \times 10^{18}$ – $3.4 \times 10^{18} \text{ kg}$ . Such a mass deficit is comparable to the expected deficit ( $\sim 3$ – $8.5 \times 10^{18} \text{ kg}$ ) from a small plume head ( $\sim 400 \text{ km}$  radius; Hill et al., 1992) that is 100–300 °C hotter than the surrounding asthenosphere.

## DISCUSSION AND PROPOSED CENOZOIC TECTONIC HISTORY

The western Cordillera is elevated because of diverse processes that date back at least to the Laramide orogeny. The Cordillera is elevated more than 1 km above sea level from the Sierra Nevada and Cascades in the west, across the Great Basin and into the Rockies and Great Plains to the east. Processes related to the horizontal subduction of the Farallon plate most likely contributed to the thick crust and high elevation of the Rocky Mountains and eastern Colorado Plateau (e.g., Dickinson and Snyder, 1978), and the region may have been at high elevation at least since Eocene time (Gregory and Chase, 1992). The period from 40 to 20 Ma represented a shift from low-angle subduction and back-arc compression to normal subduction, back-arc extension (e.g., Eaton, 1979; Zoback et al., 1981), and/or extensional collapse (e.g., Harry et al., 1993) in the Basin and Range province (Fig. 2). The localization of the back-arc and back-transform extensional setting in the Basin and Range province probably caused the lithosphere to thin relative to the more stable Colorado Plateau and Rocky Mountains.

About 16–17 Ma the Yellowstone plume emerged, and accompanying basaltic volcanism began to dominate in the northern Basin and Range. Interpretations of Neogene Cordilleran basalt composi-



**Figure 3. Current topography shown with path of Yellowstone plume. Dashed lines indicate suggested topographic expressions of starting plume head and active plume tail.**

tions vary as to whether they are consistent with a mantle plume source; some workers have suggested that they are (e.g., Leat et al., 1988; Thompson et al., 1989; Fitton et al., 1991), whereas others have found that isotopic signatures in places are more closely related to the underlying lithosphere (e.g., Lipman and Glazner, 1991; Bradshaw et al., 1993). The degree of lithospheric contamination of deep-source magmas remains controversial. A mantle plume acts more as a source of voluminous hot material than as a point source of heat, and it is likely that heat applied broadly to the lithosphere by a plume head would cause some melting of the existing mantle lithosphere as well as supply more primitive melts directly from the asthenosphere.

If the Yellowstone plume behaved like most observed plumes, then it created a broad swell centered approximately in northern Nevada. The present-day topography seems to bear the imprint of such a swell (Fig. 3), the 1 km topographic contour marking a roughly circular region centered in northwest Nevada, bounded by the slope of the Sierra Nevada to the west, the Columbia River basin to the north, the northern to southern Basin and Range transition to the south, and verging into the thickened Colorado Plateau to the east. The diameter of the observed swell in the western Cordillera is smaller ( $\sim 800 \text{ km}$ ) than the typical hot-spot swell ( $\sim 1000$ – $2000 \text{ km}$ ; e.g., Sleep, 1990; Hill et al., 1992), which may be the result of a smaller Yellowstone starting plume head caused by interaction with the subducting Juan de Fuca plate. The shape of the swell is likely to be asymmetric, because previously thinned lithosphere would create more room for plume material to pond beneath it (Fig. 2). Thus, the strongest topographic inversion would be observed in regions that were thinned prior to plume-head contact. The newly elevated Basin and Range crust was put into a strong state of extension (e.g., Crough, 1983), the northern Nevada rift forming at its center, and a new surge of broad extension began in the western part of the province during the middle Miocene. The stalk of the Yellowstone plume remained stationary as the North American plate tracked over it, introducing new basaltic volcanism to the eastern Basin and Range, as well as limited uplift resulting from plume-tail flow. The westward progression of the plate relative to the plume may have caused the broadening of the northern Basin and Range relative to the south and the most recent extension in the eastern part of the province (Pierce and Morgan, 1992). Further analysis of plume behavior in relation to the tectonic evolution of the Basin and Range

province and its high-standing margins, the Sierra Nevada and Colorado Plateau, holds great promise.

#### ACKNOWLEDGMENTS

Supported by National Science Foundation grants EAR 92-04708 and EAR 90-17667. We thank Rick Saltus (U.S. Geological Survey) for providing digital isostatic residual gravity data for use in the isostatic modeling; Mary Lou Zoback, Jill McCarthy, and Rick Saltus for reviews; and Ian Campbell, Gordon Eaton, and John Geissman for helpful comments on the manuscript.

#### REFERENCES CITED

- Armstrong, F.C., Leeman, W.P., and Malde, H.E., 1975, K-Ar dating, Quaternary and Neogene volcanic rocks of the Snake River Plain, Idaho: *American Journal of Science*, v. 275, p. 225–251.
- Beghoul, N., Barazangi, M., and Isacks, B.L., 1993, Lithospheric structure of Tibet and western North America: Mechanisms of uplift and a comparative study: *Journal of Geophysical Research*, v. 98, p. 1997–2016.
- Bradshaw, T.K., Hawkesworth, C.J., and Gallagher, K., 1993, Basaltic volcanism in the southern Basin and Range: No role for a mantle plume: *Earth and Planetary Science Letters*, v. 116, p. 45–62.
- Campbell, I.H., and Griffiths, R.W., 1990, Implications of mantle plume structure for the evolution of flood basalts: *Earth and Planetary Science Letters*, v. 99, p. 79–93.
- Catchings, R.D., and Mooney, W.D., 1989, Basin and Range crustal and upper mantle structure, northwest to central Nevada: *Journal of Geophysical Research*, v. 96, p. 6247–6267.
- Christiansen, R.L., and Lipman, P.W., 1972, Cenozoic volcanism and plate tectonic evolution of the western United States; II, Late Cenozoic: *Royal Society of London Philosophical Transactions*, v. 271, p. 234–249.
- Crough, S.T., 1983, Hotspot swells: *Annual Review of Earth and Planetary Sciences*, v. 11, p. 165–193.
- Davies, G.F., 1992, Temporal variation of the Hawaiian plume flux: *Earth and Planetary Science Letters*, v. 113, p. 277–286.
- Dickinson, W.R., and Snyder, W.S., 1978, Plate tectonics of the Laramide orogeny, in Matthews, V., III, ed., *Laramide folding associated with basement block faulting in the western United States*: Geological Society of America Memoir 151, p. 355–366.
- Duncan, R.A., and Richards, M.A., 1991, Hotspots, mantle plumes, flood basalts, and true polar wander: *Reviews of Geophysics*, v. 29, p. 31–50.
- Eaton, G.P., 1979, Regional geophysics, Cenozoic tectonics and geological resources of the Basin and Range province and adjoining regions, in Newman, G.W., and Goode, H.D., eds., *Basin and Range symposium*: Denver, Colorado, Rocky Mountain Association of Geologists, p. 11–39.
- Eaton, G.P., 1982, The Basin and Range province: Origin and tectonic significance: *Annual Review of Earth and Planetary Sciences*, v. 10, p. 409–440.
- Fitton, J.G., James, D., and Leeman, W.P., 1991, Basic magmatism associated with late Cenozoic extension in the western United States: Compositional variations in space and time: *Journal of Geophysical Research*, v. 96, p. 13,693–13,712.
- Gregory, K.M., and Chase, C.G., 1992, Tectonic significance of paleobotanically estimated climate and altitude of the late Eocene erosion surface, Colorado: *Geology*, v. 20, p. 581–585.
- Griffiths, R.W., and Campbell, I.H., 1991, On the dynamics of long-lived plume conduits in the convecting mantle: *Earth and Planetary Science Letters*, v. 103, p. 214–227.
- Hamilton, W.B., and Myers, W.B., 1966, Cenozoic tectonics of the western United States: *Reviews of Geophysics*, v. 4, p. 509–549.
- Harry, D.L., Sawyer, D.S., and Leeman, W.P., 1993, The mechanics of continental extension in western North America: Implications for the magmatic and structural evolution of the Great Basin: *Earth and Planetary Science Letters*, v. 117, p. 59–71.
- Hill, R.I., 1991, Starting plumes and continental breakup: *Earth and Planetary Science Letters*, v. 104, p. 398–416.
- Hill, R.I., Campbell, H., Davies, G.F., and Griffiths, R.W., 1992, Mantle plumes and continental tectonics: *Science*, v. 256, p. 186–192.
- Iyer, H.M., and Hitchcock, T., 1989, Upper mantle velocity structure in the continental U.S. and Canada, in Pakiser, L.C., and Mooney, W.D., eds., *Geophysical framework of the continental United States*: Geological Society of America Memoir 172, p. 681–710.
- Kohler, W., and McCarthy, J., 1990, PACE seismic refraction and wide angle reflection studies of the Colorado Plateau–Basin and Range tran- sition, Arizona: *Eos (Transactions, American Geophysical Union)*, v. 71, p. 1593.
- Lachenbruch, A.H., and Morgan, P., 1990, Continental extension, magmatism, and elevation; formal relations and rules of thumb: *Tectonophysics*, v. 174, p. 39–62.
- Leat, P.T., Thompson, R.N., Morrison, M.A., Hendry, G.D., and Dickinson, A.P., 1988, Compositionally diverse Miocene–Recent rift related magmatism in northwest Colorado: Partial melting and mixing of mafic magmas from 3 different asthenospheric and lithospheric mantle sources: *Journal of Petrology, Special Lithosphere Issue*, p. 251–377.
- Lipman, P.W., and Glazner, A.F., 1991, Introduction to middle Tertiary cordilleran volcanism: Magma sources and relations to regional tectonics: *Journal of Geophysical Research*, v. 96, p. 13,193–13,200.
- Luedke, R.G., and Smith, R.L., 1983, Map showing distribution, composition, and age of late Cenozoic volcanic centers in Idaho, western Montana, west-central South Dakota, and northwestern Wyoming: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-1091-C, scale 1:1,000,000.
- McCarthy, J., Larkin, S.P., Simpson, R.W., and Howard, K.A., 1991, Anatomy of a metamorphic core complex: Seismic refraction/wide-angle reflection profiling in southeastern California and western Arizona: *Journal of Geophysical Research*, v. 96, p. 12,259–12,291.
- Morgan, P., and Swenberg, C.A., 1985, On the Cenozoic uplift and tectonic stability of the Colorado Plateau: *Journal of Geodynamics*, v. 3, p. 39–63.
- Morgan, W.J., 1972, Deep mantle convection plume and plate motions: *American Association of Petroleum Geologists Bulletin*, v. 56, p. 203–312.
- Pierce, K.L., and Morgan, L.A., 1992, The track of the Yellowstone hot spot: Volcanism, faulting, and uplift, in Link, P.K., et al., eds., *Regional geology of eastern Idaho and western Wyoming*: Geological Society of America Memoir 179, p. 1–53.
- Prodehl, C., 1978, Crustal structure of the western United States: U.S. Geological Survey Professional Paper 1034, 74 p.
- Richards, M.A., Duncan, R.A., and Courtillot, V.E., 1989, Flood basalts and hot-spot tracks: Plume heads and tails: *Science*, v. 246, p. 103–107.
- Saltus, R.W., 1991, Gravity and heat flow constraints on Cenozoic tectonics of the western United States Cordillera [Ph.D. thesis]: Stanford, California, Stanford University, 245 p.
- Simpson, R.W., and Jachens, R.C., 1989, Gravity methods in regional studies, in Pakiser, L.C., and Mooney, W.D., eds., *Geophysical framework of the continental United States*: Geological Society of America Memoir 172, p. 35–44.
- Sleep, N.H., 1990, Hotspots and mantle plumes: Some phenomenology: *Journal of Geophysical Research*, v. 95, p. 6715–6736.
- Sparlin, M.A., Braile, L.W., and Smith, R.B., 1982, Crustal structure of the eastern Snake River Plain determined from ray trace modeling of seismic refraction data: *Journal of Geophysical Research*, v. 87, p. 2619–2633.
- Thompson, G.A., and Zoback, M.L., 1979, Regional geophysics of the Colorado Plateau: *Tectonophysics*, v. 61, p. 149–181.
- Thompson, R.N., Leat, P.T., and Humphreys, E., 1989, What is the influence of the Yellowstone mantle plume on Pliocene–Recent western USA magmatism?: *New Mexico Bureau of Mines and Mineral Resources Bulletin*, v. 131, p. 268.
- Warren, D.H., 1969, A seismic refraction survey of crustal structure in central Arizona: *Geological Society of America Bulletin*, v. 80, p. 257–282.
- Wernicke, B., 1992, Cenozoic extensional tectonics of the western U.S. Cordillera, in Burchfiel, B.C., et al., eds., *The Cordilleran orogen: Co-terminous United States*: Boulder, Colorado, Geological Society of America, *Geology of North America*, v. G-3, p. 553–581.
- Zoback, M.L., and Thompson, G.A., 1978, Basin and Range rifting in northern Nevada: Clues from a mid-Miocene rift and its subsequent offsets: *Geology*, v. 6, p. 111–116.
- Zoback, M.L., Anderson, R.E., and Thompson, G.A., 1981, Cainozoic evolution of the state of stress and style of tectonism of the Basin and Range province of the western United States: *Royal Society of London Philosophical Transactions*, ser. A, v. 300, p. 407–434.

Manuscript received June 14, 1993

Revised manuscript received September 20, 1993

Manuscript accepted September 29, 1993